




Grey-capped Tyrannulet

Grey-capped Tyrannulet E, NT

Phyllomyias griseocapilla
Piolinho-serrano

11 cm. Bright-coloured tyrannulet with contrasting yellow sides and white belly. Above olivaceous-green; coverts and flight-feathers brownish, fringed bright yellowish. These fringes form panel both on the secondaries and coverts. Fringes of tertials yellow, shaped like other *Phyllomyias*. Crown and nape greyish, contrasting with the back. Supercilium whitish, extends beyond the eye; black eye-stripe. Eye-ring white. Sides of breast and flanks are bright olivaceous-yellow, contrasting with white centre of breast and belly. Undertail coverts and vent white.



S.P. Monotypic

VOICE: Call is a simple whistle, **fyeeeee-hi**, repeated every few seconds, the last syllable lower and with a lower volume (sometimes missing altogether); another call is a series of repeated notes, **wit, wit, wit, wit**.

HABITAT: Primary and secondary forest, clearings with scattered trees, between 700 and 1,850 m.

IN AN AREA: Fairly common in the region. Penedo (Rua FF), Itatiaia NP, Mauá.

IN SE BRAZIL: Fairly common.



Grey-capped Tyrannulet

Grey-backed Tachuri E, NT

Polystictus superciliaris
Papa-moscás-de-costas-cinzentas

10 cm. Small tyrannulet of high-altitude grasslands. Above dark, olive-grey, wings dusky-brown with narrow, buffy edges to flight feathers and to the tips of wing coverts. Tail dusky-brown. Head grey with prominent white supercilium and semi-concealed, white crown-stripe; eye-ring white. Ear coverts grey, streaked white. Underparts pale buffy.



S.P. Monotypic

VOICE: Song is a fast, bubbling trill **treeheheheheheheee**, lasting about 1.5 seconds. Call is a simple, short phrase **shi-eld, shi-eld** with a ringing quality, repeated.

HABITAT: Rocky outcrops in cerrado and grassland, between 900 and 2,350 m.

IN AN AREA: Rare, only recently recorded near Itatiaia NP, at the end of the Agulhas Negras road, near Pousada Alsene.

IN SE BRAZIL: Uncommon and sparsely distributed. Serra da Canastra, Serra do Cipó, Caracá, Serra da Bocaina.



Grey-backed Tachuri

Crested Doradito



Pseudocolopteryx sclateri
Tricolino

11 cm. Inconspicuous, but easily identified, small tyrannulet of marshes. Above olive-brown with dark brown streaks; wing feathers dusky brown

with white fringes on wing coverts tips and flight feathers. Blackish crown feathers crested with semi-concealed yellowish-white crown-stripe; mask dusky. Underparts bright yellow.

S.P. Monotypic

VOICE: Call is a thin **tic**; song (not heard in the region) is said to be a four-note phrase with a squeaky quality.

HABITAT: Marshes.

IN AN AREA: Rare in the region, but frequently encountered in Resende Wetland, where it has also bred.

IN SE BRAZIL: Sparsely distributed, uncommon and erratic.



Crested Doradito

Eared Pygmy-Tyrant



Myiornis auricularis
Miudinho

7 cm. Very small bird with unmistakable head pattern. Above olivaceous-greenish; two buffy wing bars; secondaries with yellowish fringes. Crown chocolate-brown; buffy loreal area; black auricular patch and black spot below the eye; sides of neck pale greyish. Throat and upper breast whitish, streaked black. Lower breast, belly and vent bright yellow. Tail very short. Bill black.

S.P. Monotypic

VOICE: Call is a simple chipping trill, **trrrrrrrrrrr**.

HABITAT: Primary and secondary forest, forest edge.

IN AN AREA: Uncommon in the region. Penedo, Itatiaia NP, Mauá.

IN SE BRAZIL: Fairly common.



Eared Pygmy-Tyrant